arXiv:1304.4805v1 [math.DS] 17 Apr 2013

SLIDING INVARIANT AND CLASSIFICATION OF SINGULAR HOLOMORPHIC FOLIATIONS IN THE PLANE

TRUONG HONG MINH

Abstract. By introducing a new invariant called the set of slidings, we give a complete strict classification of the class of germs of non-dicritical holomorphic foliations in the plan whose Camacho-Sad indices are not rational. Moreover, we will show that, in this class, the new invariant is finitely determined. Consequently, the finite determination of the class of isoholonomy non-dicritical foliations and absolutely dicritical foliations that have the same Dulac maps are proved.

1. Introduction

The problem of classification of germs of foliations in the complex plane is stated by Thom [14]. He conjectured that the analytic type of a foliation defined in a neighborhood of a singular point is completely determined by its associated separatrix and its corresponding holonomy. R. Moussu in [12] gave a counterexample for this statement and showed that we have to consider the holonomy representation of each irreducible component of the exceptional divisor in desingularization instead of the holonomies. For this new version, the Thom's problem is proved for cuspidal type singular points [12], [4] and more generally for the quasi-homogeneous foliations [5]. However, the new statement of Thom's conjecture was denied by J.F.Mattei by computing the dimension of the space of isoholonomic deformations [10], [9]: There must be other invariants for the non quasi-homogeneous foliations. This conclusion is confirmed by the number of free coefficients in the normal forms in [6], [7] and in the hamiltonian part of the normal forms of vector field in [8]. By adding a new invariant called *set of slidings* this paper solves the problem of strict classification for the non-dicritical foliations whose Camacho-Sad indices are not rational. Here, strict classification means up to a diffeomorphism tangent to identity.

1.1. **Preliminary.**

A germ of singular foliation in $(\mathbb{C}^2, 0)$ is called *reduced* if there exists a coordinate system in which it is defined by a 1-form whose linear part is

$$\lambda_1 y dx + \lambda_2 x dy,\ \frac{\lambda_2}{\lambda_1} \notin \mathbb{Q}_{<0}.$$

Date: April 2013.

Key words and phrases. invariant of foliations, sliding invariant, classification of foliations.

$\lambda = \frac{\lambda_2}{\lambda_1}$ is called the *Camacho-Sad index* of $\mathcal{F}$. When $\lambda = 0$, the origin is called a *saddle-node* singularity, otherwise it is called *nondegenerate*. A theorem of A. Seidenberg [13, 11] said that any singular foliation $\mathcal{F}$ with isolated singularity admits a canonical *desingularization*. More precisely, there is a holomorphic map

$$\sigma : \mathcal{M} \to (\mathbb{C}^2, 0) \tag{1}$$

obtained as a composition of finite number of blowing-ups at points such that any point m of the *exceptional divisor* $\mathcal{D} := \sigma^{-1}(0)$ is either a regular point or a reduced singularity of the strict transform $\tilde{\mathcal{F}} = \sigma^*(\mathcal{F})$. An intersection of two irreducible components of $\mathcal{D}$ is called a *corner*. An irreducible component of $\mathcal{D}$ is a *dead branch* if in this component there is unique singularity of $\tilde{\mathcal{F}}$ that is a corner.

A *separatrix* of $\mathcal{F}$ is an analytical irreducible invariant curve through the origin of $\mathcal{F}$. It is well known that any germ of singular foliation $\mathcal{F}$ in $(\mathbb{C}^2, 0)$ possesses at least one separatrix [2]. When the number of separatrices is finite $\mathcal{F}$ is *non-dicritical*. Otherwise it is called *dicritical*.

Suppose that $\mathcal{F}$ is non-dicritical. Denote by $\mathrm{Sing}(\tilde{\mathcal{F}})$ the set of all singularities of the strict transform $\tilde{\mathcal{F}}$. Let D be an irreducible component of the exceptional divisor $\mathcal{D}$ then $D^* = D \setminus \mathrm{Sing}(\tilde{\mathcal{F}})$ is an leaf of $\tilde{\mathcal{F}}$. Let m be a regular point in D^* and Σ a small analytic section through m transverse to $\tilde{\mathcal{F}}$. For any loop γ in D^* based on m there is a germ of a holomorphic return map $h_\gamma : (\Sigma, m) \to (\Sigma, m)$ which only depends on the homotopy class of γ in the fundamental group $\pi_1(D^*, m)$. The map $h : \pi_1(D^*, m) \to \mathrm{Diff}(\Sigma, m)$ is called the *vanishing holonomy representation* of $\mathcal{F}$ on D. Let $\mathcal{F}'$ be a non-dicritical foliation that also admits σ as its desingularization map. Assume that $\mathrm{Sing}(\tilde{\mathcal{F}}') = \mathrm{Sing}(\tilde{\mathcal{F}})$ where $\mathrm{Sing}(\tilde{\mathcal{F}}')$ is the set of singularities of strict transform $\tilde{\mathcal{F}}'$. Denote by h'_γ in $\mathrm{Diff}(\Sigma, m)$ the vanishing holonomy representation of $\mathcal{F}$. We say that the vanishing holonomy representation of $\mathcal{F}$ and $\mathcal{F}'$ on D are conjugated if there exists $\phi \in \mathrm{Diff}(\Sigma, m)$ such that $\phi \circ h_\gamma = h'_\gamma \circ \phi$. The vanishing holonomy representation of $\mathcal{F}$ and $\mathcal{F}'$ are called conjugated if they are conjugated on every irreducible component of $\mathcal{D}$.

Notation 1. *We denote by $\mathcal{M}$ all non-dicritical foliations $\mathcal{F}$ defined on $(\mathbb{C}^2, 0)$ such that the Camacho-Sad index of $\tilde{\mathcal{F}}$ at each singularity is not rational.*

If $\mathcal{F}$ in $\mathcal{M}$ then after desingularization all the singularities of $\tilde{\mathcal{F}}$ are not saddle-node. Moreover, the Chern class of an irreducible component of divisor, that is an integer, is equal to the sum of Camacho-Sad indices of the singularities in this component [2]. Therefore, every elements in $\mathcal{M}$ after desingularization admits no dead branch in there exceptional divisors.

1.2. **Absolutely dicritical foliation.** Let σ as in (1) be a composition of finite number of blowing-ups at points. A a germ of singular holomorphic foliation $\mathcal{L}$ is said *σ-absolutely dicritical* if the strict transform $\tilde{\mathcal{L}} = \sigma^*(\mathcal{L})$ is a regular foliation and the exceptional divisor $\mathcal{D} = \sigma^{-1}(0)$ is completely transverse to the $\tilde{\mathcal{L}}$. When σ is the standard blowing-up at the origin, we called $\mathcal{L}$ is a *radial foliation.* At each corners $p = D_i \cap D_j$ of $\mathcal{D}$, the diffeomorphism from (D_i, p) to (D_j, p) that follows the leaves of $\tilde{\mathcal{L}}$ called the *Dulac map* of $\tilde{\mathcal{L}}$ at p. The existence of such foliations for any given σ is proved in [3]. In fact, in [3] the authors showed that if in each smooth component of $\mathcal{D}$ we take any two smooth curves transverse to $\mathcal{D}$ then there is always an absolutely dicritical foliation admitting them as their integral curves. We will denote by $\mathrm{Sep}(\tilde{\mathcal{F}}) \pitchfork \tilde{\mathcal{L}}$ if at any point $p \in \mathrm{Sing}(\tilde{\mathcal{F}})$ the separatrices of $\tilde{\mathcal{F}}$ through p are transverse to $\tilde{\mathcal{L}}$.

Lemma 2. *Let $\mathcal{F}$ be a non-dicritical foliation such that σ is its desingularrization map. Then there exists a σ-absolutely dicritical foliation $\mathcal{L}$ satisfying* $\mathrm{Sep}(\tilde{\mathcal{F}}) \pitchfork \tilde{\mathcal{L}}$

Proof. Denote by $L_1, \ldots, L_k$ the strict transforms of the separatrices of $\mathcal{F}$. On each component D of $\mathcal{D}$ that does not contain any singularity of $\tilde{\mathcal{F}}$ except the corners we take a smooth curve L_{k+j} transverse to D. Then we have the set of curve $\{L_1, \ldots, L_n\}$ such that each component of $\mathcal{D}$ is transverse to at least one curve L_i. Denote by $p_i = L_i \cap \mathcal{D}$. By [3], for each i there exists an absolutely dicritical foliation $\mathcal{L}_i$ defined by a 1-form ω_i verifying that L_i is transverse to $\tilde{\mathcal{L}}_i$. Choose a local chart (x_i, y_i) at p_i such that $\mathcal{D} = \{x_i = 0\}, L_i = \{y_i = 0\}$ and

$$\sigma^*\omega_i(x_i, y_i) = x_i^{m_i} d(x_i + y_i) + h.o.t,$$

where h.o.t stands for higher order term. Write down ω_i in the local chart (x_j, y_j)

$$\sigma^*\omega_i(x_j, y_j) = x_j^{m_j} d(a_{ij}x_j + b_{ij}y_j) + h.o.t. \tag{2}$$

Because $\tilde{\mathcal{L}}_i$ is transverse to $\mathcal{D}$, we have $b_{ij} \neq 0$. We define $a_{ii} = b_{ii} = 1$. There always exits a vector $(c_1, \ldots, c_n) \in \mathbb{C}^n$ such that for $j = 1, \ldots, n$

$$a_j = \sum_{i=1}^{n} c_i a_{ij} \neq 0 \ \text{ and } \ b_j = \sum_{i=1}^{n} c_i b_{ij} \neq 0.$$

Denote by $\omega_0 = \sum c_i \omega_i$. Then, in the local chart (x_j, y_j), we have

$$\sigma^*\omega_0(x_j, y_j) = x_j^{m_j} d(a_j x_j + b_j y_j) + h.o.t. \text{ where } a_j \neq 0, b_j \neq 0.$$

Because ω_0 and ω_i, $i = 1, \ldots, n$, have the same multiplicity on each component of $\mathcal{D}$, they have the same vanishing order. Since each component of $\mathcal{D}$ contains at least a point p_i and the strict transform $\tilde{\mathcal{L}}$ of the foliation $\mathcal{L}$ defined by ω_0 is transverse to $\mathcal{D}$ in each neighborhood of each p_i, $\tilde{\mathcal{L}}$ is generically transverse to $\mathcal{D}$. By [3], $\mathcal{L}$ is absolutely dicritical and it is the one we need. □

1.3. **Slidings of foliations.** Consider first a nondegenerate reduced foliation $\mathcal{F}$ in $(\mathbb{C}^2, 0)$. By [11], there exists a coordinate system in which $\mathcal{F}$ is defined by

$$\lambda y(1 + A(x, y))dx + xdy, \ \lambda \notin \mathbb{Q}_{\leq 0},$$

where $A(0, 0) = 0$. Let $\mathcal{L}$ be a germ regular foliation whose invariant curve through the origin (we call it the separatrix of $\mathcal{L}$) is transverse to the two separatrices of $\mathcal{F}$ that are S_1 and S_2. Then we have the following Lemma, whose proof is straightforward.

Lemma 3. *The tangent curve of $\mathcal{F}$ and $\mathcal{L}$, denote by $T(\mathcal{F}, \mathcal{L})$, is a smooth curve transverse to the two separatrices of $\mathcal{F}$. Moreover, if the separatrix of $\mathcal{L}$ is tangent to $\{x - cy = 0\}$ then $T(\mathcal{F}, \mathcal{L})$ is tangent to $\{x + c\lambda y = 0\}$.*

After a standard blowing-up σ_1 at the origin, the strict transform $\tilde{T}(\mathcal{F}, \mathcal{L})$ of $T(\mathcal{F}, \mathcal{L})$ is transverse to $\tilde{\mathcal{F}}$ and cut $D_1 = \sigma_1^{-1}(0)$ at p. We denote by $D_1^* = D_1 \setminus \operatorname{Sing}(\sigma_1^*(\mathcal{F}))$ and $\tilde{h} : \pi_1(D_1^*, p) \to \operatorname{Diff}(\tilde{T}(\mathcal{F}, \mathcal{L}), p)$ the vanishing holonomy representation of $\mathcal{F}$. We choose a generator γ for $\pi_1(D_1^*, p) \cong \mathbb{Z}$. Then σ_1 induces

$$h_\gamma = \sigma_1 \circ \tilde{h}(\gamma) \circ \sigma_1^{-1} \in \operatorname{Diff}(T(\mathcal{F}, \mathcal{L}), 0).$$

We call h_γ the *holonomy on the tangent curve* $T(\mathcal{F}, \mathcal{L})$. Denote by π_{S_1} and π_{S_2} the projection by the leaves of $\mathcal{L}$ from $T(\mathcal{F}, \mathcal{L})$ to S_1 and S_2 respectively.

Definition 4. *The sliding of a reduced foliation $\mathcal{F}$ and a regular foliation $\mathcal{L}$ on S_1 (resp. S_2) is the diffeomorphism*

$$g_{S_1}(\mathcal{F}, \mathcal{L}) = \pi_{S_1 *}(h_\gamma) = \pi_{S_1} \circ h_\gamma \circ \pi_{S_1}^{-1}$$
$$\big(\textit{resp. } g_{S_2}(\mathcal{F}, \mathcal{L}) = \pi_{S_2 *}(h_\gamma) = \pi_{S_2} \circ h_\gamma \circ \pi_{S_2}^{-1}\big).$$

Let $d : S_1 \to S_2$ the Dulac map of $\mathcal{L}$ (Section 1.2). Since $d = \pi_y \circ \pi_x^{-1}$, it is obvious that

$$g_{S_2}(\mathcal{F}, \mathcal{L}) = d_* \left(g_{S_1}(\mathcal{F}, \mathcal{L})\right). \tag{3}$$

Now let $\mathcal{F}$ be a non-dicritical foliation such that after desingularization by the map σ all singularities of $\sigma^*(\mathcal{F}) = \tilde{\mathcal{F}}$ are nondegenerate. By Lemma 2 there exists a σ-absolutely dicritical foliation $\mathcal{L}_0$ such that $\operatorname{Sep}(\tilde{\mathcal{F}}) \pitchfork \tilde{\mathcal{L}}_0$.

Notation 5. *We denote by $\mathcal{R}(\mathcal{L}_0)$ the set of all σ-absolutely dicritical foliations $\mathcal{L}$ satisfying the two following properties*

- *$\tilde{\mathcal{L}}$ and $\tilde{\mathcal{L}}_0$ has the same the Dulac maps at any corners of $\mathcal{D}$.*
- *At each singularities of $\tilde{\mathcal{F}}$, the invariant curves of $\tilde{\mathcal{L}}$ and $\tilde{\mathcal{L}}_0$ are tangent.*

Let $\mathcal{L}$ in $\mathcal{R}(\mathcal{L}_0)$ and D be a irreducible component of $\mathcal{D}$. Suppose that $p_1, \ldots, p_m$ the singularities of $\tilde{\mathcal{F}}$ on D. Then we denote by

$$S_D(\tilde{\mathcal{F}}, \tilde{\mathcal{L}}) = \{g_{D,p_1}(\tilde{\mathcal{F}}, \tilde{\mathcal{L}}), \ldots, g_{D,p_m}(\tilde{\mathcal{F}}, \tilde{\mathcal{L}})\},$$

where $g_{D,p_i}(\tilde{\mathcal{F}}, \tilde{\mathcal{L}})$ is the sliding of $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{L}}$ in a neighborhood of p_i.


S_2
$\mathcal{L}$
$\mathcal{F}$
$T(\mathcal{F},\mathcal{L})$
$g_{S_2}(\mathcal{F},\mathcal{L})$
h_γ
$g_{S_1}(\mathcal{F},\mathcal{L})$
O
S_1

FIGURE 1. Sliding of $\mathcal{F}$ and $\mathcal{L}$

Definition 6. *The sliding of* $\mathcal{F}$ *and* $\mathcal{L}$ *is*

$$S(\mathcal{F},\mathcal{L}) = \cup_{D\in \mathrm{Comp}(\mathcal{D})} S_D(\tilde{\mathcal{F}},\tilde{\mathcal{L}}),$$

where Comp$(\mathcal{D})$ *is the set of all irreducible components of* $\mathcal{D}$*. The set of slidings of* $\mathcal{F}$ *relative to direction* $\mathcal{L}_0$ *is the set*

$$\mathcal{S}_0(\mathcal{F}) = \cup_{\mathcal{L}\in\mathcal{R}(\mathcal{L}_0)} S(\mathcal{F},\mathcal{L}).$$

We will prove in Corollary 12 that $\mathcal{S}_0(\mathcal{F})$ is an invariant of $\mathcal{F}$: If $\mathcal{F}$ and $\mathcal{F}'$ are conjugated by Φ then $S(\mathcal{F},\mathcal{L}) = \tilde{\Phi}_{|\mathcal{D}} \circ S(\mathcal{F}',\Phi_*\mathcal{L}) \circ \tilde{\Phi}^{-1}_{|\mathcal{D}}$. Under some conditions for $\mathcal{F}$ and $\mathcal{F}'$ (Theorem 8), we will have $\tilde{\Phi}_{|\mathcal{D}} = \mathrm{Id}$. Therefore $S(\mathcal{F},\mathcal{L}) = S(\mathcal{F}',\Phi_*\mathcal{L})$. Consequently, $\mathcal{S}_0(\mathcal{F}) = \mathcal{S}_0(\mathcal{F}')$.

Remark 7. *For each singularity* p *of* $\tilde{\mathcal{F}}$ *that is a corner; i.e,* $p = D_i \cap D_j$*, there are two slidings* $g_{D_i,p}(\tilde{\mathcal{F}},\tilde{\mathcal{L}})$ *and* $g_{D_j,p}(\tilde{\mathcal{F}},\tilde{\mathcal{L}})$*. However, by* (3)*,* $g_{D_j,p}(\tilde{\mathcal{F}},\tilde{\mathcal{L}})$ *is completely determined by* $g_{D_i,p}(\tilde{\mathcal{F}},\tilde{\mathcal{L}})$ *and the Dulac map of* $\tilde{\mathcal{L}}$ *at* p*.*

This invariant is named "sliding" because it gives an obstruction for the construction of local conjugacy of two foliation that fixes the points in exceptional divisor (Corollary 12).

Although $S(\mathcal{F},\mathcal{L})$ *is the set of local diffeomorphism, it also contains the additional informations through the "global" foliation* $\mathcal{L}$*.*

1.4. **Statement of the main results.** Let $\mathcal{F},\mathcal{F}'\in\mathcal{M}$. We say that their *strict separatrices are tangent*, denote $\operatorname{Sep}(\tilde{\mathcal{F}})\,/\!\!/\operatorname{Sep}(\tilde{\mathcal{F}}')$, if they have the same desingularization map and the same set of singularities, moreover, at each singularity which is not a corner of divisor the separatrices of $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$ are tangent. If $\operatorname{Sep}(\tilde{\mathcal{F}})\,/\!\!/\operatorname{Sep}(\tilde{\mathcal{F}}')$ and $\mathcal{L}_0$ is an absolutely dicritical foliation satisfying $\operatorname{Sep}(\tilde{\mathcal{F}})\pitchfork\tilde{\mathcal{L}}_0$ then $\operatorname{Sep}(\tilde{\mathcal{F}})\pitchfork\tilde{\mathcal{L}}_0'$. We also denote by $\operatorname{CS}(\tilde{\mathcal{F}})$ the set of Camacho-Sad indices of $\tilde{\mathcal{F}}$ at all singularities. Let denote by $\operatorname{CS}(\tilde{\mathcal{F}})=\operatorname{CS}(\tilde{\mathcal{F}}')$ if at each singularity, $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$ have the same Camacho-Sad index

Theorem 8. *Let* $\mathcal{F}$ *and* $\mathcal{F}'$ *be two foliations in the class* $\mathcal{M}$ *(see Notation 1) such that* $\operatorname{Sep}(\tilde{\mathcal{F}})\,/\!\!/\operatorname{Sep}(\tilde{\mathcal{F}}')$*. Suppose that* $\mathcal{L}_0$ *is an absolutely dicritical foliation satisfying* $\operatorname{Sep}(\tilde{\mathcal{F}})\pitchfork\tilde{\mathcal{L}}_0$*. Let* $\mathcal{R}(\mathcal{L}_0)$ *as in Notation 5 and* $\mathcal{S}_0(\mathcal{F})$ *and* $\mathcal{S}_0(\mathcal{F}')$ *the corresponding set of slidings. Then the tree following statements are equivalent*

(i) $\mathcal{F}$ *and* $\mathcal{F}'$ *are strictly analytically conjugated.*
(ii) *Their vanishing holonomy representation are strictly analytically conjugated,* $\operatorname{CS}(\tilde{\mathcal{F}})=\operatorname{CS}(\tilde{\mathcal{F}}')$ *and* $\mathcal{S}_0(\mathcal{F})=\mathcal{S}_0(\mathcal{F}')$
(iii) *Their vanishing holonomy representation are strictly analytically conjugated,* $\operatorname{CS}(\tilde{\mathcal{F}})=\operatorname{CS}(\tilde{\mathcal{F}}')$ *and* $\mathcal{S}_0(\mathcal{F})\cap\mathcal{S}_0(\mathcal{F}')\neq\emptyset$

Here, a strict conjugacy means a conjugacy tangent to identity. We will prove that the sliding of foliations are finitely determined:

Theorem 9. *Let* $\mathcal{F}$ *be a non-dicritical foliation without saddle-node singularity after desingularization. There exists a natural* N *such that if there is a non-dicritical foliation* $\mathcal{F}'$ *satisfying the following conditions:*

(i) $\mathcal{F}$ *and* $\mathcal{F}'$ *have the same set of singularities after desingularization and at neighborhood of each singularity* $\tilde{\mathcal{F}}$ *and* $\tilde{\mathcal{F}}'$ *are locally strictly analytically conjugated,*
(ii) *There exist* $\mathcal{L},\mathcal{L}'$ *in* $\mathcal{R}(\mathcal{L}_0)$ *such that* $J^N(S(\mathcal{F},\mathcal{L}))=J^N(S(\mathcal{F}',\mathcal{L}'))$*,*

then there exists $\mathcal{L}''$ *such that* $\mathcal{L}''$ *is strictly conjugated with* $\mathcal{L}$ *and* $S(\mathcal{F},\mathcal{L}'')=S(\mathcal{F}',\mathcal{L}')$*.*

Where $J^N(S(\mathcal{F},\mathcal{L}))=J^N(S(\mathcal{F}',\mathcal{L}'))$ means $J^N(g_{D,p}(\tilde{\mathcal{F}},\tilde{\mathcal{L}}))=J^N(g_{D,p}(\tilde{\mathcal{F}}',\tilde{\mathcal{L}}'))$ for all $g_{D,p}(\tilde{\mathcal{F}},\tilde{\mathcal{L}})$ in $S(\mathcal{F},\mathcal{L})$, $g_{D,p}(\tilde{\mathcal{F}}',\tilde{\mathcal{L}}')$ in $S(\mathcal{F}',\mathcal{L}')$ and $J^N(g_{D,p}(\tilde{\mathcal{F}},\tilde{\mathcal{L}}))$ stands for the regular part of degree N in the Taylor expansion of $g_{D,p}(\tilde{\mathcal{F}},\tilde{\mathcal{L}})$.

These two theorems also give two corollaries about finite determination of the class of isoholonomy non-dicritical foliations and absolutely dicritical foliations that have the same Dulac maps (see Corollary 20 and 22).

This paper is organized as follows: In section 2, local conjugacy of the pair $(\mathcal{F},\mathcal{L})$ will be proved. We prove Theorem 8 in Section 3. Section 4 devotes to prove Theorem 9 and two Corollaries of finite determination of class of isoholonomy non-dicritical foliations and absolutely dicritical foliations that have the same Dulac maps.

2. Local conjugacy of the pair $(\mathcal{F},\mathcal{L})$

Let $\mathcal{F}$, $\mathcal{F}'$ be two germs of nondegenerate reduced foliations in $(\mathbb{C}^2,0)$. Denote by S_1, S_2 and S_1', S_2' the separatrices of $\mathcal{F}$ and $\mathcal{F}'$ respectively. Let $\mathcal{L}$ and $\mathcal{L}'$ be two germs of regular foliations such that their separatrices L and L' are transverse to the two separatrices of $\mathcal{F}$ and $\mathcal{F}'$ respectively. Suppose that Φ is a diffeomorphism conjugating $(\mathcal{F},\mathcal{L})$ and $(\mathcal{F}',\mathcal{L}')$ then the restriction of Φ on the tangent curves commutes with the holonomies on $T(\mathcal{F},\mathcal{L})$ and $T(\mathcal{F}',\mathcal{L}')$ of $\mathcal{F}$ and $\mathcal{F}'$. The converse is also true:

Proposition 10. *Suppose that $\mathcal{F}$ and $\mathcal{F}'$ have the same Camacho-Sad index. If $\phi : T(\mathcal{F},\mathcal{L}) \to T(\mathcal{F}',\mathcal{L}')$ is a diffeomorphism commuting with the holonomies of $\mathcal{F}$ and $\mathcal{F}'$ then ϕ extends to a diffeomorphism Φ of $(\mathbb{C}^2,0)$ sending $(\mathcal{F},\mathcal{L})$ to $(\mathcal{F}',\mathcal{L}')$. Moreover, if we require that Φ sends S_1, S_2 to S_1', S_2' respectively then this extension is unique.*

Proof. By Lemma 2, the curves S_1, S_2, L, $T(\mathcal{F},\mathcal{L})$ (resp. S_1', S_2', L', $T(\mathcal{F}',\mathcal{L}')$) are four transverse smooth curves. It is well known that there exist two radial foliations $\mathcal{R}$ and $\mathcal{R}'$ such that S_1, S_2, L, $T(\mathcal{F},\mathcal{L})$ and S_1', S_2', L, $T(\mathcal{F}',\mathcal{L}')$ are the invariant curves of $\mathcal{R}$ and $\mathcal{R}'$ respectively. After a blowing-up at the origin, denote by p_1, p_2, p_L, p_T (resp. p_1', p_2', p_L', p_T') the intersection of strict transforms of S_1, S_2, L, $T(\mathcal{F},\mathcal{L})$ (resp. S_1', S_2', L, $T(\mathcal{F}',\mathcal{L}')$) with $\mathbb{CP}^1$. Let ϕ_1 in $\mathrm{Aut}(\mathbb{CP}^1)$ that sends p_1, p_2, p_L to p_1', p_2', p_L' respectively. Then $\phi_1(p_T) = p_T'$ by Lemma 2. Using the path lifting method after a blowing-up [11], ϕ can extends to a diffeomorphism Φ_1 of $(\mathbb{C}^2,0)$ sending $(\mathcal{F},\mathcal{R})$ to $(\mathcal{F}',\mathcal{R}')$. Denote by $\mathcal{L}_0 = \Phi_{1*}^{-1}(\mathcal{L}')$. Because Φ_1^{-1} sends L' and $T(\mathcal{F}',\mathcal{L}')$ to L and $T(\mathcal{F},\mathcal{L})$ respectively, L is also the separatrix of $\mathcal{L}_0$ and $T(\mathcal{F},\mathcal{L}_0) = T(\mathcal{F},\mathcal{L})$. We denote by T the tangent curve $T(\mathcal{F},\mathcal{L}_0)$. The proof is reduced to show that there exists a diffeomorphism fixing points in T sending $(\mathcal{F},\mathcal{L})$ to $(\mathcal{F},\mathcal{L}_0)$. Choose a system of coordinates (x,y) such that $\mathcal{L}_0$ is defined by $f_0 = x+y$ and $\mathcal{F}$ is defined by a 1-form

$$\omega(x,y) = \lambda y(1 + A(x,y))dx + xdy,\ \lambda \notin \mathbb{Q}_{\leq 0}.$$

Then T is defined by

$$\tau(x,y) = x - \lambda y(1 + A(x,y)) = 0.$$

We claim that there exist a natural $n \geq 2$ and $h \in \mathbb{C}\{x,y\}$ such that $\mathcal{L}$ is defined by

$$f(x,y) = (1 + \tau^n(x,y)h(x,y))\,(x+y).$$

Indeed, assume that $\mathcal{L}$ is defined by

$$\bar{f}(x,y) = u(x,y)(x+y).$$

There exists a diffemorphism $g \in \mathbb{C}\{y\}$ such that

$$g\Big(u\Big(\lambda y(1+A), y\Big).\Big(\lambda y(1+A)+y\Big)\Big) = \lambda y(1+A)+y.$$

It is equivalent to

$$\big(g \circ \bar{f} - (x+y)\big)_{|\tau=0} = 0.$$

Then $\mathcal{L}$ is also defined by $f = g \circ \bar{f}$ and there exist a natural n and a function h satisfying $n \geq 1$ and $h_{|\tau=0} \not\equiv 0$ such that

$$g \circ \bar{f}(x,y) = (1 + \tau^n(x,y)h(x,y))\,(x+y).$$

Moreover

$$\begin{aligned} df \wedge \omega &= \tau(x,y)(\ldots) + n(x+y)h(x,y)\tau^{n-1} d\tau \wedge \omega \\ &= \tau(x,y)(\ldots) + n(x+y)h(x,y)\tau^{n-1}(x + \lambda^2 y + \ldots)dx \wedge dy. \end{aligned}$$

The fact that $\big(df \wedge \omega\big)_{|\tau=0} \equiv 0$ and $\lambda \neq 0, -1$ forces to $n \geq 2$.
Let

$$X = x\frac{\partial}{\partial x} - \lambda y(1 + A(x,y))\frac{\partial}{\partial y}$$

tangent to $\mathcal{F}$. Now we will show that there exists $\alpha \in \mathbb{C}\{x,y\}$ such that the diffeomorphism $\exp[\tau^{n-1}\alpha]X$ satisfies

$$(x+y) \circ \exp[\tau^{n-1}\alpha]X(x,y) = \sum_{i\geq 0} \frac{\tau^{i(n-1)}\alpha^i}{i!}\mathrm{ad}^i_X(x+y) = f(x,y), \tag{4}$$

where ad_X is the adjoint representation. Since

$$\sum_{i\geq 0} \frac{\tau^{i(n-1)}\alpha^i}{i!}\mathrm{ad}^i_X(x+y) = x + y + \tau^n\alpha + \frac{n}{2}\tau^{2n-2}\alpha^2 X(\tau) + \tau^{2n-1}(\ldots),$$

(4) becomes

$$\alpha + \frac{n}{2}\tau^{n-2}\alpha^2 X(\tau) + \tau^{n-1}(\ldots) = (x+y)h(x,y).$$

Hence, the existence of α comes from the implicit function theorem.

Now we will prove the uniqueness of Φ. In fact, we only need to show that if there exists a diffeomorphism Ψ that sends $(\mathcal{F}, \mathcal{L}_0)$ to itself, preserves the two separatrices of $\mathcal{F}$ and fixes the points of T then $\Psi = \mathrm{Id}$. Since $\Psi_{|T} = \mathrm{Id}$, Ψ sends every leaf of $\mathcal{F}$ into itself. By [1], there exists $\beta \in \mathbb{C}\{x,y\}$ such that

$$\Psi = \exp[\beta]X.$$

Because $\mathcal{L}_0$ is defined by the function $x+y$ and Ψ fixes points in T, we get

$$(x+y) \circ \exp[\beta]X = x+y. \tag{5}$$

Decompose β into the homogeneous terms

$$\beta = \beta_0 + \beta_1 + \beta_2 + \ldots = \beta_0 + \bar{\beta}.$$

Since $\mathrm{ad}_X^i(x) = x$ and $\mathrm{ad}_X^i(y) = (-\lambda)^i y + c_i$ for all i, where $c_i \in (x,y)^2$, we have

$$\begin{aligned}(x+y)\circ\exp[\beta]X &= \sum_{i=0}^{\infty}\frac{\beta_0^i}{i!}x + \sum_{i=0}^{\infty}\frac{\beta_0^i}{i!}((-\lambda)^i y) + h.o.t\\ &= \exp(\beta_0)x + \exp(-\lambda\beta_0)y + h.o.t.\end{aligned}$$

So (5) leads to

$$\exp(\beta_0) = \exp(-\lambda\beta_0) = 1.$$

Hence,

$$x\circ\exp[\beta_0]X = \sum_{i=0}^{\infty}\frac{\beta_0^i}{i!}x = \exp(\beta_0)x = x, \tag{6}$$

$$y\circ\exp[\beta_0]X = \sum_{i=0}^{\infty}\frac{\beta_0^i}{i!}\left((-\lambda)^i y + c_i\right) = \exp(-\lambda\beta_0)y + c = y + c, \tag{7}$$

where $c \in (x,y)^2$. We claim that

$$\exp[\beta]X = \exp[\beta_0]X\circ\exp[\bar\beta]X. \tag{8}$$

Indeed, for any $h \in \mathbb{C}\{x,y\}$ we have

$$\begin{aligned}h\circ\exp[\beta_0]X\circ\exp[\bar\beta]X &= \left(\sum_{i=0}^{\infty}\frac{\beta_0^i}{i!}\mathrm{ad}_X^i(h)\right)\circ\exp[\bar\beta]X = \sum_{j=0}^{\infty}\frac{\bar\beta^j}{j!}\mathrm{ad}_X^j\left(\sum_{i=0}^{\infty}\frac{\beta_0^i}{i!}\mathrm{ad}_X^i(h)\right)\\ &= \sum_{k=0}^{\infty}\sum_{i+j=k}\frac{\bar\beta^j\beta_0^i}{j!i!}\mathrm{ad}_X^k(h) = \sum_{k=0}^{\infty}\frac{(\bar\beta+\beta_0)^k}{k!}\mathrm{ad}_X^k(h) = h\circ\exp[\beta]X.\end{aligned}$$

We write $\mathrm{ad}_X^i(y+c) = (-\lambda)^i y + d_i$ where $d_i \in (x,y)^2$, by (6), (7), (8) we get

$$\begin{aligned}(x+y)\circ\exp[\beta]X &= x\circ\exp[\beta_0]X\circ\exp[\bar\beta]X + y\circ\exp[\beta_0]X\circ\exp[\bar\beta]X\\ &= x\circ\exp[\bar\beta]X + (y+c)\circ\exp[\bar\beta]X\\ &= \sum_{i=0}^{\infty}\frac{\bar\beta^i}{i!}x + \sum_{i=0}^{\infty}\frac{\bar\beta^i}{i!}\mathrm{ad}_X^i(y+c)\\ &= \exp(\bar\beta)x + \exp(-\lambda\bar\beta)y + \sum_{i=0}^{\infty}\frac{\bar\beta^i}{i!}d_i\\ &= x\prod_{i=1}^{\infty}\exp(\beta_i) + y\prod_{i=1}^{\infty}\exp(-\lambda\beta_i) + \sum_{i=0}^{\infty}\frac{\bar\beta^i}{i!}d_i. \qquad (9)\end{aligned}$$

We will prove $\bar\beta = 0$ by induction. From (9) we have

$$(x+y)\circ\exp[\beta]X = x(1+\beta_1) + y(1-\lambda\beta_1) + h.o.t.$$

So (5) forces $\beta_1 = 0$. Suppose that $\beta_1 = \ldots = \beta_{k-1} = 0$, we have

$$(x+y)\circ\exp[\beta]X = x(1+\beta_k) + y(1-\lambda\beta_k) + h.o.t.$$

Then (5) again leads to $\beta_k = 0$ and consequently $\beta = \beta_0$. It implies that

$$\Psi = \exp[\beta_0]X = (x, y + c).$$

Finally, (5) again gives $c = 0$. So $\Psi = \mathrm{Id}$. ☐

Corollary 11. *Suppose that $\mathcal{F}$ and $\mathcal{F}'$ are two nondegenerate reduced foliations that are analytically conjugated. Let $\mathcal{L}$ and $\mathcal{L}'$ be two regular foliation that are transverse to the two separatrices of $\mathcal{F}$ and $\mathcal{F}'$ respectively. Then there exits a diffeomorphism that sends $(\mathcal{F}, \mathcal{L})$ to $(\mathcal{F}', \mathcal{L}')$.*

Proof. Let Ψ be the conjugacy of $\mathcal{F}$ and $\mathcal{F}'$. Denote by $T' = \Psi(T(\mathcal{F}, \mathcal{L}))$. Then the restriction $\Psi_{|T(\mathcal{F},\mathcal{L})}$ commutes with the holonomies of $\mathcal{F}$ on $T(\mathcal{F}, \mathcal{L})$ and $\mathcal{F}'$ on T'. Moreover by the holonomy transport, the holonomies of $\mathcal{F}'$ on T' and $T(\mathcal{F}', \mathcal{L}')$ are conjugated. Hence, the holonomies of $\mathcal{F}$ on $T(\mathcal{F}, \mathcal{L})$ and $\mathcal{F}'$ on $T(\mathcal{F}', \mathcal{L}')$ are conjugated. By Proposition 10 there exits a diffeomorphism that sends $(\mathcal{F}, \mathcal{L})$ to $(\mathcal{F}', \mathcal{L}')$ ☐

By projecting on S_1 and S_2 the holonomies defined on $T(\mathcal{F}, \mathcal{L})$ and $T(\mathcal{F}', \mathcal{L}')$ respectively, we can obtain

Corollary 12. *If Φ is a diffeomorphism conjugating $(\mathcal{F}, \mathcal{L})$ and $(\mathcal{F}', \mathcal{L}')$, then*

$$\Phi_{|S_1} \circ g_{S_1}(\mathcal{F}, \mathcal{L}) = g_{S_1'}(\mathcal{F}', \mathcal{L}') \circ \Phi_{|S_1}.$$

Reciprocally, if $\mathrm{CS}(\mathcal{F}) = \mathrm{CS}(\mathcal{F}')$ and $\phi : S_1 \to S_1'$ is a diffeomorphism satisfying

$$\phi \circ g_{S_1}(\mathcal{F}, \mathcal{L}) = g_{S_1'}(\mathcal{F}', \mathcal{L}') \circ \phi$$

then ϕ uniquely extends to a diffeomorphism Φ of $(\mathbb{C}^2, 0)$ sending $(\mathcal{F}, \mathcal{L})$ to $(\mathcal{F}', \mathcal{L}')$.

Proof. Because Φ conjugates $(\mathcal{F}, \mathcal{L})$ and $(\mathcal{F}', \mathcal{L}')$, the restriction $\Phi_{|T(\mathcal{F},\mathcal{L})}$ commutes with the holonomies h_γ and h'_γ of $\mathcal{F}$ and $\mathcal{F}'$. Denote by π_{S_1} (resp. $\pi_{S_1'}$) the projection by the leaves of $\mathcal{L}$ (resp. $\mathcal{L}'$) from $T(\mathcal{F}, \mathcal{L})$ (resp. $T(\mathcal{F}', \mathcal{L}')$) to S_1 (resp. S_1'). Since Φ sends $(\mathcal{F}, \mathcal{L})$ to $(\mathcal{F}', \mathcal{L}')$, we have

$$\pi_{S_1'} \circ \Phi_{|T(\mathcal{F},\mathcal{L})} = \Phi_{|S_1} \circ \pi_{S_1}.$$

Therefore

$$\begin{aligned}
\Phi_{|S_1} \circ g_{S_1}(\mathcal{F}, \mathcal{L}) &= \Phi_{|S_1} \circ \pi_{S_1} \circ h_\gamma \circ \pi_{S_1}^{-1} = \pi_{S_1'} \circ \Phi_{|T(\mathcal{F},\mathcal{L})} \circ h_\gamma \circ \pi_{S_1}^{-1} \\
&= \pi_{S_1'} \circ h'_\gamma \circ \Phi_{|T(\mathcal{F},\mathcal{L})} \circ \pi_{S_1}^{-1} = g_{S_1'}(\mathcal{F}', \mathcal{L}') \circ \pi_{S_1'} \circ \Phi_{|T(\mathcal{F},\mathcal{L})} \circ \pi_{S_1}^{-1} \\
&= g_{S_1'}(\mathcal{F}', \mathcal{L}') \circ \Phi_{|S_1}.
\end{aligned}$$

Reciprocally, if $\phi : S_1 \to S_1'$ is a diffeomorphism commuting with the slidings of $\mathcal{F}$ and $\mathcal{F}'$. Denote by $\psi = \pi_{S_1'}^{-1} \circ \phi \circ \pi_{S_1}$ then

$$\begin{aligned}
\psi \circ h_\gamma &= \pi_{S_1'}^{-1} \circ \phi \circ \pi_{S_1} \circ h_\gamma = \pi_{S_1'}^{-1} \circ \phi \circ g_{S_1}(\mathcal{F}, \mathcal{L}) \circ \pi_{S_1} \\
&= \pi_{S_1'}^{-1} \circ g_{S_1'}(\mathcal{F}', \mathcal{L}') \circ \phi \circ \pi_{S_1} = h'_\gamma \circ \pi_{S_1'}^{-1} \circ \phi \circ \pi_{S_1} = h'_\gamma \circ \psi.
\end{aligned}$$

By Proposition 10, ψ uniquely extends to a diffeomorphism Φ that sends $(\mathcal{F},\mathcal{L})$ to $(\mathcal{F}',\mathcal{L}')$. ☐

Remark 13. *In particular, if in Corollary 12 we have that* $S_1 = S_1'$ *and* $g_{S_1}(\mathcal{F},\mathcal{L}) = g_{S_1'}(\mathcal{F}',\mathcal{L}')$ *then there exists a diffeomorphism sending* $(\mathcal{F},\mathcal{L})$ *to* $(\mathcal{F}',\mathcal{L}')$ *and fixing points in* S_1.

3. Strict classification of foliations in $\mathcal{M}$

This whole section devotes to prove Theorem 8.

Proof of theorem 8. The direction ((ii)⇒(iii)) is obvious.

((i)⇒(ii)) Let Φ be the strict conjugacy and $\tilde{\Phi} : (\mathcal{M},\mathcal{D}) \to (\mathcal{M},\mathcal{D})$ be its lifting by σ. Suppose that a non-corner point m of $\mathcal{D}$ is a fixed point of $\tilde{\Phi}$. Then the linear map $D\tilde{\Phi}(m)$ has two eigenvalues. One is corresponding to the direction of divisor. We denote by $v(\tilde{\Phi})(m)$ the other and define $v(\tilde{\Phi})(m) = 1$ for each corner m.

Lemma 14. $\tilde{\Phi}_{|\mathcal{D}} = \mathrm{Id}$ *so* $v(\tilde{\Phi})$ *is a function defined on* $\mathcal{D}$ *and moreover* $v(\tilde{\Phi}) \equiv 1$.

Proof. Denote by σ_1 the standard blowing-up at origin of $(\mathbb{C}^2, 0)$

$$\sigma_1 : (\mathcal{M}_1, D_1) \to (\mathbb{C}^2, 0).$$

On D_1, we use the two standard chart $(x,\bar{y})$ and $(\bar{x}, y)$ together with the transition functions $\bar{x} = \bar{y}^{-1}$, $y = x\bar{y}$. Suppose that

$$\Phi(x,y) = (x + \alpha(x,y), y + \beta(x,y)),\ \alpha,\beta \in (x,y)^2.$$

Then in the coordinate system $(x,\bar{y})$ we have

$$\begin{aligned}\Phi_1(x,\bar{y}) = \sigma_1^*\Phi(x,\bar{y}) &= \left(x + \alpha(x,x\bar{y}), \frac{x\bar{y} + \beta(x,x\bar{y})}{x + \alpha(x,x\bar{y})}\right)\\ &= (x(1+\ldots), y + x(\beta_0 + \ldots)),\end{aligned}$$

where $\beta = \beta_0 x^2 + \ldots$. Therefore $\Phi_1 : (\mathcal{M}_1, \mathcal{D}_1) \to (\mathcal{M}_1, D_1)$ fixes points in D_1 and $v(\Phi_1) \equiv 1$. Let p be a singularity of $\sigma_1^*\mathcal{F}$ on D_1. We will show that $D\Phi_1(p) = \mathrm{Id}$ and apply the inductive hypothesis for Φ_1 in a neighborhood of p . Indeed, let σ_2 be the blowing-up at p and $D_2 = \sigma_2^{-1}(p)$. Denote by S_p and S_p' all invariant curves of $\sigma_1^*\mathcal{F}$ and $\sigma_1^*\mathcal{F}'$ through p. Because every elements in $\mathcal{M}$ after desingularization admits no dead branch in there exceptional divisors, D_2 is not a dead component. Therefore there is at least one irreducible component of S_p and S_p' that are not tangent to D_1. Because $\Phi_{1*}(S_p) = S_p'$, $D\phi_1(p)$ has an eigenvector different from the direction of D_1. So $D\Phi_1(p)$ has two eigenvectors. Since both of their eigenvalues are 1, we have $D\Phi_1(p) = \mathrm{Id}$. ☐

Now let $\mathcal{L} \in \mathcal{R}_0$ and denote by $\mathcal{L}' = \Phi_*(\mathcal{L})$. Since $\tilde{\Phi}_{|\mathcal{D}} = \mathrm{Id}$, the strict transforms $\tilde{\mathcal{L}}$ and $\tilde{\mathcal{L}}'$ have the same Dulac maps. Moreover, because $\mathrm{Sep}(\tilde{\mathcal{F}}) \,/\!\!/\, \mathrm{Sep}(\tilde{\mathcal{F}}')$, at each singularity p of $\tilde{\mathcal{F}}$, $D\tilde{\Phi}(p)$ has two eigenvectors. As $v(\tilde{\Phi}) \equiv 1$ and $\tilde{\Phi}_{|\mathcal{D}} = \mathrm{Id}$ we have $D\tilde{\Phi}(p) = \mathrm{Id}$. Therefore the invariant curves of $\tilde{\mathcal{L}}$ and $\tilde{\mathcal{L}}'$ through p are tangent. This gives $\mathcal{L}' = \Phi_*(\mathcal{L}) \in \mathcal{R}(\mathcal{L}_0)$. Because $\tilde{\Phi}$ fixes points in $\mathcal{D}$, by Corollary 12 the identity map commutes with the slides of $\mathcal{F}$ and $\mathcal{F}'$. It leads to $S(\mathcal{F}, \mathcal{L}) = S(\mathcal{F}', \mathcal{L}')$. Consequently, $\mathcal{S}_0(\mathcal{F}) = \mathcal{S}_0(\mathcal{F}')$. Moreover, the vanishing holonomy representation of $\mathcal{F}$ and $\mathcal{F}'$ are conjugated by $\tilde{\Phi}$. Since $v(\tilde{\Phi}) \equiv 1$ the conjugacy are strict.

((iii)⇒(i)) Suppose that $\mathcal{L}$, $\mathcal{L}' \in \mathcal{R}_0$ satisfy $S(\mathcal{F}, \mathcal{L}) = S(\mathcal{F}', \mathcal{L}')$. By Corollary 12, at each singularity p_i, $i \in \{1, \ldots, k\}$, of $\tilde{\mathcal{F}}$ there exists a neighborhood U_i of p_i and a local conjugacy

$$\Phi_i : (\tilde{\mathcal{F}}, \tilde{\mathcal{L}})_{|U_i} \to (\tilde{\mathcal{F}}', \tilde{\mathcal{L}}')_{|U_i}$$

such that $\Phi_{i|\mathcal{D}\cap U_i} = \mathrm{Id}$. Let U_0 be a neighborhood of $\mathcal{D} \setminus \cup_{i=1}^k U_i$ such that U_0 does not contain any singularity of $\tilde{\mathcal{F}}$. Note that U_0 is not connected and the restriction of $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$ on U_0 are regular. The strict conjugacy of the vanishing holonomy representations can be extended by path lifting method to the conjugacy

$$\Phi_0 : (\tilde{\mathcal{F}}, \tilde{\mathcal{L}})_{|U_0} \to (\tilde{\mathcal{F}}', \tilde{\mathcal{L}}')_{|U_0},$$

satisfying that the second eigenvalue function $v(\Phi_0) \equiv 1$. We will show that on each intersection $V_i = U_i \cap U_0$, Φ_i and Φ_0 are coincident. Denote by

$$\Psi_i = \Phi_{i|V_i}^{-1} \circ \Phi_{0|V_i} : (\tilde{\mathcal{F}}, \tilde{\mathcal{L}})_{|V_i} \to (\tilde{\mathcal{F}}, \tilde{\mathcal{L}})_{|V_i}.$$

We claim that $v(\Psi_i) \equiv 1$ on $\mathcal{D} \cap V_i$. Let p, q in $V_i \cap \mathcal{D}$. Denote by l_p and l_q the invariant curves of $\tilde{\mathcal{L}}$ through p and q respectively. As the two maps $\Psi_{i|l_p}$ and $\Psi_{i|l_q}$ are conjugated by the holonomy transport, we have $v(\Psi_i)(p) = v(\Psi_i)(q)$. Consequently, $v(\Psi_i)$ is constant on V_i. Since $v(\Phi_0) \equiv 1$, it follows that $v(\Phi_i)$ is constant on $V_i \cap \mathcal{D}$. Therefore, $v(\Phi_i)$ is constant on $U_i \cap \mathcal{D}$. Moreover, at the singularity p_i, $D\Phi_i(p_i)$ has three eigenvectors corresponding to the directions of the divisor and the directions of invariant curves of $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{L}}$ through p_i. Since $D\Phi_i(p_i)$ has also one eigenvalue 1 corresponding to the directions of the divisor, we have $D\Phi_i(p_i) = \mathrm{Id}$. This gives $v(\Phi_i) \equiv 1$ and consequently $v(\Psi_i) \equiv 1$.

Now at each point $p \in V_i \cap \mathcal{D}$, the map $\Psi_{i|l_p}$ commutes with the holonomy of $\tilde{\mathcal{F}}$ around p_i. Since the Camacho-Sad index λ_i at p_i is not rational, Lemma 15 below says that $\Psi_{i|l_p} = \mathrm{Id}$ and so $\Psi_i = \mathrm{Id}$. Hence we can paste all Φ_i together and the strict conjugacy we need is the projection of this diffeomorphism on $(\mathbb{C}^2, 0)$ by σ. □

Lemma 15. *Let $h \in \mathrm{Diff}(\mathbb{C}, 0)$ such that $h'(0) = \exp(2\pi i\lambda)$ where $\lambda \notin \mathbb{Q}$. If $\psi \in \mathrm{Diff}(\mathbb{C}, 0)$ satisfying $\psi'(0) = 1$ and $\psi \circ h = h \circ \psi$ then $\psi = \mathrm{Id}$*

Proof. Since $|h'(0)| = |\exp(2\pi i\lambda)| \neq 1$, there is a formal diffeomorphism ϕ such that $\phi \circ h \circ \phi^{-1}(z) = \exp(2\pi i\lambda)z$. Denote by $\tilde{\psi} = \phi \circ \psi \circ \phi^{-1}$, then $\tilde{\psi}'(0) = 1$ and $\tilde{\psi}(\exp(2\pi i\lambda)z) = \exp(2\pi i\lambda)\tilde{\psi}$. The proof is reduced to show that $\tilde{\psi} = \mathrm{Id}$. Suppose that $\tilde{\psi}(z) = z + \sum_{j=2}^{\infty} a_j z^j$. Then

$$\tilde{\psi}(\exp(2\pi i\lambda)z) = \exp(2\pi i\lambda)z + \sum_{j=2}^{\infty} a_j \exp(2j\pi i\lambda)z^j,$$

and

$$\exp(2\pi i\lambda)\tilde{\psi}(z) = \exp(2\pi i\lambda)z + \sum_{j=2}^{\infty} a_j \exp(2\pi i\lambda)z^j.$$

Since $\lambda \notin \mathbb{Q}$, it forces $a_j = 0$ for all $j \geq 2$. Hence $\tilde{\psi} = \mathrm{Id}$. □

4. Finite determinacy

Let S be a germ of smooth curve in $(\mathbb{C}^2, 0)$. Denote by $\mathfrak{m}^n(S)$ the set of all holomorphic functions on S whose vanishing orders at the origin are at least n. We also denote by $\Sigma(S)$ the set of all germs of smooth curves having the same desingularization map and having the same singularities as S after desingularization.

Proposition 16. *Let S be a germ of curve in $(\mathbb{C}^2, 0)$ and $S_1, \ldots, S_k$ be its irreducible components. Suppose that $\sigma : (\mathcal{M}, \mathcal{D}) \to (\mathbb{C}^2, 0)$ be a finite composition of blowing-ups such that all the transformed curves $\sigma^* S_i = \tilde{S}_i$ are smooth. Then there exists a natural N such that if $f_i \in \mathfrak{m}^N(\tilde{S}_i)$, $i = 1, \ldots, k$ then there exists $F \in \mathbb{C}\{x, y\}$ such that $F \circ \sigma_{|\tilde{S}_i} = f_i$. Moreover N can be chosen the same for all elements in $\Sigma(S)$*

Proof. We first consider the statement when S is irreducible. If S is smooth then $\tilde{S}$ is diffeomorphic to S. So we can suppose that S is singular. Denote by $p = \tilde{S} \cap \mathcal{D}$. Choose a coordinate system (x_p, y_p) in a neighborhood of p such that $\tilde{S} = \{y_p = 0\}$ and $\mathcal{D} = \{x_p = 0\}$. Then σ^{-1} is defined by

$$x_p = \frac{\alpha(x, y)}{\beta(x, y)} \text{ and } y_p = \frac{\mu(x, y)}{\nu(x, y)},$$

where $\alpha, \beta, \mu, \nu \in \mathbb{C}\{x, y\}$, $\gcd(\alpha, \beta) = 1$ and $\gcd(\mu, \nu) = 1$. So we have

$$\frac{\alpha \circ \sigma(x_p, y_p)}{\beta \circ \sigma(x_p, y_p)} = x_p \tag{10}$$

Therefore, there exist a natural k and a holomorphic function h such that

$$\alpha \circ \sigma(x_p, y_p) = x_p^{k+1} h(x_p, y_p),\ \beta \circ \sigma(x_p, y_p) = x_p^k h(x_p, y_p), \tag{11}$$

where $x_p \nmid h$. We claim that h is a unit. Indeed, if $h(0, 0) = 0$, denote by $\tilde{L}$ the curve $\{h(x_p, y_p) = 0\}$. Let L be a curve defined in $(\mathbb{C}^2, 0)$ such that $\sigma^*(L) = \tilde{L}$. Let $\{\tilde{h}(x, y) = 0\}$ be the equation of L. By (11), $\tilde{h}|\alpha$ and $\tilde{h}|\beta$.

It contradicts $\gcd(\alpha, \beta) = 1$. Now denote by $u(x_p) = h(x_p, 0)$ that is a unit, we have

$$\alpha \circ \sigma(x_p, 0) = u(x_p)x_p^{k+1},\ \beta \circ \sigma(x_p, 0) = u(x_p)x_p^k. \tag{12}$$

For each $m \geq (k-1)(k+1)$ there exists $j \in \{0, \ldots, k-1\}$ such that $k|(m - j(k+1))$. Thus

$$m = ik + j(k+1),\ i, j \in \mathbb{N}.$$

So (12) implies that if a holomorphic function $f(x_p)$ satisfying $x_p^{(k-1)(k+1)}|f(x_p)$ then there exists a holomorphic function $F(x, y)$ such that $F \circ \sigma(x_p, 0) = f(x_p)$. Consequently

$$\mathfrak{m}^{(k-1)(k+1)}(\tilde{S}) \subset \sigma^*\mathbb{C}\{x, y\}_{|\tilde{S}}. \tag{13}$$

In the general case, suppose that S_i is defined by $\{g_i = 0\}$. If $f_i \in \mathfrak{m}^N(\tilde{S}_i)$, $i = 1, \ldots, k$, with N is big enough, there exist F_i, $i = 1, \ldots, k$, such that $F_i \circ \sigma_{|\tilde{S}_i} = f_i$. We will find a holomorphic function F such that $F_{|S_i} = F_{i|S_i}$ for all $i = 1, \ldots, k$. It is reduced to show that there exits a natural M such that the following morphism Θ is surjective

$$\frac{(x, y)^M}{(g_1) \cap \ldots \cap (g_k) \cap (x, y)^M} \to \frac{(x, y)^M}{(g_1) \cap (x, y)^M} \oplus \ldots \oplus \frac{(x, y)^M}{(g_k) \cap (x, y)^M}.$$

Indeed, by Hilbert's Nullstellensatz, there exits a natural M_1 such that

$$(x, y)^{M_1} \subset (g_i, g_j) \tag{14}$$

for all $1 \leq i < j \leq k$. We will show that for all $i = 1, \ldots, k$, $j = 0, \ldots, (k-1)N_1$ the elements $e_{ij} = (0, \ldots, \overline{x^j y^{(k-1)M_1 - j}}, \ldots, 0)$, where $\overline{x^j y^{(k-1)M_1 - j}}$ is in the i^{th} position, are in $\mathrm{Im}\Theta$ and then $M = (k-1)M_1$. We decompose

$$x^j y^{(k-1)M_1 - j} = \prod_{\substack{l=1,\ldots,k \\ l \neq i}} x^{j_l} y^{M_1 - j_l},$$

where $0 \leq j_l \leq M_1$. By (14), there exist $a_{il}, b_{il} \in \mathbb{C}\{x, y\}$ such that $a_{il}g_i + b_{il}g_l = x^{j_l}y^{M_1 - j_l}$. It implies that

$$e_{ij} = \Theta\left(\prod_{\substack{l=1,\ldots,k \\ l \neq i}} \left(x^{j_l}y^{M_1 - j_l} - a_{il}g_i\right)\right) \in \mathrm{Im}\Theta$$

Now we will show that N can be chosen the same for all elements of $\Sigma(S)$. In the case S is irreducible, let S' in $\Sigma(S)$ and $y_p = s(x_p)$ be the equation of $\sigma^*(S') = \tilde{S}'$. We also have

$$\alpha \circ \sigma(x_p, s(x_p)) = v(x_p)x_p^{k+1},\ \beta \circ \sigma(x_p, s(x_p)) = v(x_p)x_p^k,$$

where $v(x_p) = h(x_p, s(x_p))$ that is a unit. Consequently, (13) holds. In the general case, it is sufficient to show that M_1 in (14) can be chosen the same for all elements of $\Sigma(g_i)$, $i = 1, \ldots, k$. Let M_{ij} be the smallest natural satisfying

$$(x, y)^{M_{ij}} \subset (g_i, g_j).$$

we claim that

$$M_{ij} \leq \mathrm{I}(g_i, g_j) = \dim_{\mathbb{C}} \frac{\mathbb{C}\{x, y\}}{(g_i, g_j)}.$$

Indeed, there exists $x^l y^{M_{ij}-1-l} \notin (g_i, g_j)$. Let P_m, $m = 1, \ldots, M_{ij}$, be a sequence of monomials such that $P_1 = 1$, $P_{M_{ij}} = x^l y^{M_{ij}-1-l}$ and either $P_{m+1} = x \cdot P_m$ or $P_{m+1} = y \cdot P_m$. Since $P_m | P_{M_{ij}}$ we have $P_m \notin (g_i, g_j)$ for all $m = 1, \ldots, M_{ij}$. We will show that $\{P_1, \ldots, P_{M_{ij}}\}$ are independent in the vector space $\frac{\mathbb{C}\{x,y\}}{(g_i,g_j)}$ over $\mathbb{C}$. Suppose that

$$c_1 P_1 + \ldots + c_{M_{ij}} P_{M_{ij}} \in (g_i, g_j).$$

If there exists $c_m \neq 0$. Let m_0 be the smallest natural such that $c_{m_0} \neq 0$. Then

$$c_{m_0} P_{m_0} + \ldots + c_{M_{ij}} P_{M_{ij}} = P_{m_0}(c_{m_0} + \ldots) \in (g_i, g_j).$$

It implies that P_{m_0} in (g_i, g_j) and it is a contradiction.

Now, it is well known that the intersection number $\mathrm{I}(g_i, g_j)$ is a topological invariant. It means that if two curves $\{g_i \cdot g_j = 0\}$ and $\{g'_i \cdot g'_j = 0\}$ are topologically conjugated then $\mathrm{I}(g_i, g_j) = \mathrm{I}(g'_i, g'_j)$. Consequently, M_1 can be chosen as $max_{1 \leq i < j \leq k} \mathrm{I}(g_i, g_j)$ that doesn't depend on the elements of $\Sigma(g_i)$, $i = 1, \ldots, k$. □

Proof of Theorem 9. Suppose that $\tilde{T}(\mathcal{F}, \mathcal{L}) = \cup T_i$, $\tilde{T}'(\mathcal{F}', \mathcal{L}') = \cup T'_i$ where T_i and T'_i are irreducible components of $T(\mathcal{F}, \mathcal{L})$ and $T(\mathcal{F}', \mathcal{L}')$. Then the singularities of $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$: $p_i = T_i \cap \mathcal{D} = T'_i \cap \mathcal{D}$. Denote by $h_{i\gamma}$ the holonomy of $\tilde{\mathcal{F}}$ on T_i. Consider at a neighborhood of p_i.

We first suppose that p_i is not a corner. Denote by D the irreducible component of $\mathcal{D}$ through p_i. Because $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$ are strictly conjugated in a neighborhood of p_i, by Corollaries 11 and 12, there exists a diffeomorphism ψ_i in $\mathrm{Diff}(D, p_i)$ tangent to identity such that

$$\psi_i \circ g_{D,p_i}(\tilde{\mathcal{F}}, \tilde{\mathcal{L}}) = g_{D,p_i}(\tilde{\mathcal{F}}', \tilde{\mathcal{L}}') \circ \psi_i. \tag{15}$$

Let π_D (resp. π'_D) be the projection from T_i (resp. T'_i) to D that follows the leaves of $\tilde{\mathcal{L}}$ (resp. $\tilde{\mathcal{L}}'$). Denote by

$$\phi_i = \pi_D^{-1} \circ \psi_i^{-1} \circ \pi_D. \tag{16}$$

Since $J^N(S(\mathcal{F}, \mathcal{L})) = J^N(S(\mathcal{F}', \mathcal{L}'))$, ϕ_i is tangent to identity map at order at least N.

In the case p_i is a corner. Let D be one of two irreducible components of $\mathcal{D}$ trough p_i and define ϕ_i as above.

Lemma 17. *Suppose that there exists a diffeomorphism Φ such that the lifting $\sigma^*(\Phi) = \tilde{\Phi}$ satisfies*

- $\tilde{\Phi}_{|\mathcal{D}} = \mathrm{Id}$,

- $\tilde{\Phi}_{|T_i} = \phi_i$,
- $T(\mathcal{F}, \Phi_* \mathcal{L}) = T(\mathcal{F}, \mathcal{L})$.

Then $\mathcal{L}'' = \Phi_*(\mathcal{L})$ *satisfies* $S(\mathcal{F}, \mathcal{L}'') = S(\mathcal{F}', \mathcal{L}')$.

Proof. We first consider in the neighborhood U_i of a non-corner singularity p_i. We denote D, π_D, π'_D as above. Let π''_D be the projection following the leaves of $\tilde{\mathcal{L}}''$ from T_i to D then

$$\pi''_D = \pi_D \circ \phi_i^{-1}.$$

We have

$$\begin{aligned} g_{D,p_i}(\tilde{\mathcal{F}}, \tilde{\mathcal{L}}'') &= \pi''_D \circ h_{i\gamma} \circ \pi''^{-1}_D = \pi_D \circ \phi_i^{-1} \circ h_{i\gamma} \circ \phi_i \circ \pi_D^{-1} \\ &= \psi_i \circ \pi_D \circ h_{i\gamma} \circ \pi_D^{-1} \circ \psi_i^{-1} = \psi_i \circ g_{D,p_i}(\tilde{\mathcal{F}}, \tilde{\mathcal{L}}) \circ \psi_i^{-1} \\ &= g_{D,p_i}(\tilde{\mathcal{F}}', \tilde{\mathcal{L}}'). \end{aligned}$$

If p_i is a corner, $p_i = D \cap D'$, we also have

$$g_{D,p_i}(\tilde{\mathcal{F}}, \tilde{\mathcal{L}}'') = g_{D,p_i}(\tilde{\mathcal{F}}', \tilde{\mathcal{L}}').$$

Since $\tilde{\Phi}_{|\mathcal{D}} = \mathrm{Id}$ the Dulac maps of $\tilde{\mathcal{L}}''$ and $\tilde{\mathcal{L}}'$ in a neighborhood of p_i are the same. So Remark 7 leads to

$$g_{D',p_i}(\tilde{\mathcal{F}}, \tilde{\mathcal{L}}'') = g_{D',p_i}(\tilde{\mathcal{F}}', \tilde{\mathcal{L}}').$$

□

Now we will prove the existence of Φ in Lemma 17 for N is big enough. Suppose that $\mathcal{F}$ and $\mathcal{L}$ are respectively defined by

$$\begin{aligned} \omega &= a(x,y)dx + b(x,y)dy, \\ \omega_{\mathcal{L}} &= c(x,y)dx + d(x,y)dy. \end{aligned}$$

Then the tangent curve $T = T(\mathcal{F}, \mathcal{L})$ is defined by

$$q(x,y) = da - cb = 0.$$

Let $X_q = \frac{\partial q}{\partial y}\frac{\partial}{\partial x} - \frac{\partial q}{\partial x}\frac{\partial}{\partial y}$ be a vector field tangent to T and $\tilde{X}_q$ be its lifting by σ. By the implicit theorem, if N is big enough, there exists f_i defined on T_i such that

$$\exp[f_i]\left(\tilde{X}_q|_{T_i}\right) = \phi_i.$$

Using Lemma 16, by choosing a big enough N, there exists $f \in \mathbb{C}\{x,y\}$ such that

$$\exp[f \circ \sigma]\tilde{X}_q|_{T_i} = \phi_i.$$

For each $\Phi = (\Phi_1, \Phi_2) \in \mathrm{Diff}(\mathbb{C}^2, 0)$, denote by

$$< \Phi > = \frac{\omega_{\mathcal{L}} \wedge \Phi^* \omega_{\mathcal{L}}}{dx \wedge dy} = c\left(c \circ \Phi \frac{\partial \Phi_1}{\partial y} + d \circ \Phi \frac{\partial \Phi_2}{\partial y}\right) - d\left(c \circ \Phi \frac{\partial \Phi_1}{\partial x} + d \circ \Phi \frac{\partial \Phi_2}{\partial x}\right).$$

It is easy to see that $T(\mathcal{F}, \Phi_* \mathcal{L}) = T$ if and only if $q| < \Phi >$. For each holomorphic function f, we denote by

$$\Phi_f = \exp[f] X_q.$$

Lemma 18 below says that there exists a holomorphic function u such that Φ_{f-uq} satisfies Lemma 17 for N is big enough. Moreover, by Proposition 16, we can chose N that does not depend on $\mathcal{L}$. $\square$

Lemma 18. *If N is big enough, for all f in $(x,y)^N$ there exists a holomorphic function u such that $q| < \Phi_{f-uq} >$.*

Proof. We have

$$\begin{aligned}\frac{\partial}{\partial x} x\circ\Phi_{f-uq}\Big|_{\{q=0\}} &= \frac{\partial}{\partial x}\sum_{i=0}^{\infty}\frac{(f-uq)^i}{i!}\mathrm{ad}^i_{X_q}(x)\Big|_{\{q=0\}}\\ &= \frac{\partial}{\partial x} x\circ\Phi_f\Big|_{\{q=0\}} - u.\frac{\partial q}{\partial x}.\sum_{i=1}^{\infty}\frac{f^{i-1}}{(i-1)!}\mathrm{ad}^i_{X_q}(x)\Big|_{\{q=0\}}\\ &= \frac{\partial}{\partial x} x\circ\Phi_f\Big|_{\{q=0\}} - u.\frac{\partial q}{\partial x}.\frac{\partial q}{\partial y}\circ\Phi_f\Big|_{\{q=0\}}.\end{aligned}$$

Similarly,

$$\begin{aligned}\frac{\partial}{\partial y} x\circ\Phi_{f-uq}\Big|_{\{q=0\}} &= \frac{\partial}{\partial y} x\circ\Phi_f\Big|_{\{q=0\}} - u\cdot\frac{\partial q}{\partial y}\cdot\frac{\partial q}{\partial y}\circ\Phi_f\Big|_{\{q=0\}},\\ \frac{\partial}{\partial x} y\circ\Phi_{f-uq}\Big|_{\{q=0\}} &= \frac{\partial}{\partial x} y\circ\Phi_f\Big|_{\{q=0\}} + u\cdot\frac{\partial q}{\partial x}\cdot\frac{\partial q}{\partial x}\circ\Phi_f\Big|_{\{q=0\}},\\ \frac{\partial}{\partial y} y\circ\Phi_{f-uq}\Big|_{\{q=0\}} &= \frac{\partial}{\partial y} y\circ\Phi_f\Big|_{\{q=0\}} + u\cdot\frac{\partial q}{\partial y}\cdot\frac{\partial q}{\partial x}\circ\Phi_f\Big|_{\{q=0\}}.\end{aligned}$$

It implies that

$$\begin{aligned}&< \Phi_{f-uq} > |_{\{q=0\}} = < \Phi_f > |_{\{q=0\}}\\ &\qquad - u\cdot\left(c\frac{\partial q}{\partial y} - d\frac{\partial q}{\partial x}\right)\cdot\left(\left(c\frac{\partial q}{\partial y} - d\frac{\partial q}{\partial x}\right)\circ\Phi_f\right)\Big|_{\{q=0\}}.\end{aligned}\tag{17}$$

Denote by $h = c\frac{\partial q}{\partial y} - d\frac{\partial q}{\partial x}$. Then $\{h=0\}$ is the tangent curve of $\mathcal{L}$ and the foliation defined by level sets of q. Since at each singularity p_i of $\tilde{\mathcal{F}}$, the irreducible component T_i of T is transverse to $\tilde{\mathcal{L}}$, the irreducible components of the strict transform of $\{h=0\}$ at p_i are also transverse to T_i. It implies that $(q,h)=1$ and the two curves $\{q\cdot h=0\}$ and $\{q\cdot(h\circ\Phi_f)=0\}$ are topologically conjugated. By Hilbert's Nullstellensatz and the proof of Proposition 16 there exists a natural M such that $(x,y)^M\subset(q,h)$ and $(x,y)^M\subset(q,h\circ\Phi_f)$. It implies that $(x,y)^{2M}\subset(q,h\cdot(h\circ\Phi_f))$. So for $< \Phi_f >$ in $(x,y)^{2M}$, by (17) we can choose $u\in\mathbb{C}\{x,y\}$ such that $q| < \Phi_{f-uq} >$. $\square$

Remark 19. *If we replace the condition "$\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$ are locally analytically conjugated" in Theorem 9 by the condition "$\mathcal{F}$ and $\mathcal{F}'$ are in $\mathcal{M}$" then the conclusion in Theorem 9 becomes: "For all natural $M\geq N$ there exists $\mathcal{L}''_M$ such that $J^M(S(\mathcal{F},\mathcal{L}''_M)) = J^M(S(\mathcal{F}',\mathcal{L}'))$". Indeed, in that case, because* *the Camacho-Sad indices are not rational, $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$ are locally formally conjugated. So we can choose ψ in* (15) *such that*

$$J^M(\psi_i \circ g_{D,p_i}(\tilde{\mathcal{F}},\tilde{\mathcal{L}}) \circ \psi_i^{-1}) = J^M(g_{D,p_i}(\tilde{\mathcal{F}}',\tilde{\mathcal{L}}')).$$

Corollary 20. *Let $\mathcal{F} \in \mathcal{M}$ defined by a 1-form ω then there exists a natural N such that if $\mathcal{F}' \in \mathcal{M}$ defined by a 1-form ω' satisfying that $J^N\omega = J^N\omega'$ and the vanishing holonomy representations of $\mathcal{F}$ and $\mathcal{F}'$ are strictly analytically conjugated then $\mathcal{F}$ and $\mathcal{F}'$ are strictly analytically conjugated.*

Proof. Let $\mathcal{L} \in \mathcal{R}_0$ then $J^{m(N)}S(\mathcal{F},\mathcal{L}) = J^{m(N)}S(\mathcal{F}',\mathcal{L})$ where $m(M)$ is an increasing function on N and $m(N) \to \infty$ when $N \to \infty$. By Theorem 9 if N is big enough there exists $\mathcal{L}'' \in \mathcal{R}_0$ such that $S(\mathcal{F},\mathcal{L}'') = S(\mathcal{F}',\mathcal{L})$. By Theorem 8, $\mathcal{F}$ and $\mathcal{F}'$ are strictly analytically conjugated. □

Remark 21. *This Corollary is consistent with the result of J. F. Mattei in* [9] *which says that the dimension of moduli space of the equisingular unfolding of a foliation is finite. Note that the vanishing holonomy representations of two foliations that are jointed by a unfolding are conjugated but the converse is not true.*

Corollary 22. *Let $\mathcal{L}$ be a σ-absolutely dicritical foliation defined by 1-form ω. There exists a natural N such that if $\mathcal{L}'$ is a σ-absolutely dicritical foliation defined by ω' satisfying that $J^N\omega = J^N\omega'$ and the Dulac maps of $\tilde{\mathcal{L}}$ and $\tilde{\mathcal{L}}'$ are the same then $\mathcal{L}$ and $\mathcal{L}'$ are strictly analytically conjugated.*

Proof. Suppose that $\mathcal{D} = \cup_{i=1,\ldots,k} D_i$ where D_i is an irreducible component of $\mathcal{D}$. We take a pair of irreducible functions f_i and g_i for each $i = 1,\ldots,k$ such that the curve $C_i = \{f_i = 0\}$ and $C_i' = \{g_i = 0\}$ satisfy the following properties:

1. The strict transforms $\tilde{C}_i$ and $\tilde{C}_i'$ cut D_i at two different points p_i, q_i, respectively, such that no one of them is a corner.
2. $\tilde{C}_i$, $\tilde{C}_i'$ are smooth and transverse to the invariant curve of $\tilde{\mathcal{L}}$ through p_i, q_i respectively.

Because $[\mathbb{C} : \mathbb{Q}]$ is an infinite field extension, there exists $(\lambda_1,\lambda_2,\ldots,\lambda_k) \in \mathbb{C}^k$ such that

$$\sum_{i=1}^{k} c_i\lambda_i \notin \mathbb{Q},\ \forall (c_1,\ldots,c_k) \in \mathbb{Q}^k \setminus \{(0,\ldots,0)\}.$$

Now, let us consider the non-dicritical foliation $\mathcal{F}$ defined by the 1-form

$$\omega_0 = \prod_{i=1}^{k}(f_ig_i)\cdot\left(\sum_{i=1}^{k}\left(\lambda_i\frac{df_i}{f_i} + \frac{dg_i}{g_i}\right)\right).$$

Then $\mathcal{F}$ admits σ as its desingularization map and the singularities of the strict transform $\tilde{\mathcal{F}}$ are the corners of $\mathcal{D}$ and p_i, q_i, $i = 1\ldots,k$. We claim that at each singularity, the Camacho-Sad index of $\tilde{\mathcal{F}}$ is not rational. Indeed, denote by m_{ij} the multiplicity of $f_i \circ \sigma$ and $g_i \circ \sigma$ on D_j. At the corner $p_{ij} =$

$D_i \cap D_j$, we take a coordinates (x, y) such that $D_i = \{x = 0\}, D_j = \{y = 0\}$. In this coordinate system, we can write $\sigma^*\omega_0$ as

$$\sigma^*\omega_0 = u(x,y)x^{2\sum_{l=1}^{k} m_{li}} y^{2\sum_{l=1}^{k} m_{lj}} \sum_{l=1}^{k} \left((\lambda_l + 1)m_{li}\frac{dx}{x} + (\lambda_l + 1)m_{lj}\frac{dy}{y} + \alpha_{ij} \right),$$

where $u(x, y)$ is a unit and α_{ij} is a holomophic form. So the Camacho-Sad index of $\tilde{\mathcal{F}}$ at p_{ij} is

$$\mathrm{I}(p_{ij}) = \frac{\sum_{l=1}^{k}(\lambda_l + 1)m_{lj}}{\sum_{l=1}^{k}(\lambda_l + 1)m_{li}} \notin \mathbb{Q}.$$

Similarly, the Camacho-Sad indices of $\tilde{\mathcal{F}}$ at p_i and q_i, respectively, are

$$\mathrm{I}(p_i) = \frac{\sum_{l=1}^{k}(\lambda_l + 1)m_{li}}{\sum_{l=1}^{k}\lambda_l} \notin \mathbb{Q},\ \mathrm{I}(q_i) = \frac{\sum_{l=1}^{k}(\lambda_l + 1)m_{li}}{k} \notin \mathbb{Q}.$$

Now if $J^N\omega = J^N\omega'$ then $J^{m(N)}S(\mathcal{F}, \mathcal{L}) = J^{m(N)}S(\mathcal{F}, \mathcal{L}')$ where $m(N)$ is an increasing function on N and $m(N) \to \infty$ when $N \to \infty$. Moreover if N is big enough the invariant curves of $\tilde{\mathcal{L}}$ and $\tilde{\mathcal{L}}'$ trough the singularities of $\tilde{\mathcal{F}}$ are tangent. By using Theorem 9 for $\mathcal{F}' = \mathcal{F}$, there exists a foliation $\mathcal{L}''$ strictly conjugated with $\mathcal{L}$ such that the two couple $(\mathcal{F}, \mathcal{L}'')$ and $(\mathcal{F}, \mathcal{L}')$ are strictly conjugated. Consequently, $\mathcal{L}$ and $\mathcal{L}'$ are strictly conjugated. □

Acknowledgements

This paper is based on the main part of my doctoral thesis in Institute of Mathematics of Toulouse, France. I would like to thank my advisors,Yohann Genzmer and Emmanuel Paul, for having guided me so well all over these years, Jean-François Mattei for useful discussions and suggestions about the problem.

References

[1] Berthier, M.; Cerveau, D.; Meziani, R. - *Transformations isotropes des germes de feuilletages holomorphes*, J. Math. Pures Appl. **9** 78 (1999), no. 7, 701–722.

[2] Camacho, C.; Sad, P. - *Invariant varieties through singularities of holomorphic vector fields*, Ann. of Math. **115**, no. 3 (1982) 579–595.

[3] Cano, F; Corral, N - *Absolutely dicritical foliations*, Int. Math. Res. Not. **8** (2011) 1926–1934.

[4] Cerveau, D.; Moussu, R. - *Groupes d'automorphismes de* $(\mathbb{C}, 0)$ *et équations différentielles* $ydy + \cdots = 0$.Bull. Soc. Math. France **116** (1988), no. 4, 459–488 (1989).

[5] Genzmer, Y. - *Analytical and formal classifications of quasi-homogeneous foliations in* $(\mathbb{C}^2, 0)$., Differential Equations **245** (2008), no. 6, 1656–1680.

[6] Genzmer, Y.; Paul, E. - *Normal forms of foliations and curves defined by a function with a generic tangent cone.* Mosc. Math. J. **11** (2011), no. 1, 41–72, 181.

[7] Genzmer, Y.; Paul, E. - *Moduli spaces for topologically quasi-homogeneous functions.*

[8] L. Ortiz-Bobadilla; E. Rosales-González; S. M. Voronin - *Thom's problem for the orbital analytic classification of degenerate singular points of holomorphic vector fields on the plane.* Dokl. Math. **82** no. 2, (2010), 759–761.

[9] Mattei, J.-F. - *Modules de feuilletages holomorphes singuliers. I. Équisingularité*, Invent. Math. **103** (1991), no. 2, 297–325.

[10] Mattei, J.-F. - *Quasi-homogénéité et équiréductibilité de feuilletages holomorphes en dimension deux*, Astérisque No. 261 (2000), xix, 253–276.

[11] Mattei, J.-F.; Moussu, R. - *Holonomy and first integrals*, Ann. Sci. École Norm. Sup. (4) **13** (1980), no. 4, 469–523.

[12] Moussu, R. - *Holonomie évenescente des équations différentielles dégénerées transverses*, Singularities and dynamical systems (Iráklion, 1983), 161–173, North-Holland Math. Stud., 103, North-Holland, Amsterdam, 1985.

[13] Seidenberg, A. - *Reduction of singularities of the differential equation* $Ady = Bdx$. , Amer. J. Math. **90** (1968) 248–269. .

[14] Żoładek, H. - *The monodromy group*, Birkhäuser Verlag, Basel, 2006. xii+580 pp.

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR5219, UNIVERSITÉ TOULOUSE 3
E-mail address: `hong-minh.truong@math.univ-toulouse.fr`